

Date _____	TEPEZZA Referral Form	
-------------------	------------------------------	--

PATIENT INFORMATION:

Last Name:	First Name:	Home Phone:	Mobile:
Home Address:		City:	State: Zip:
Delivery Address:		City:	State: Zip:
Date of Birth:	Gender:	Language:	Email:
Caregiver Name:		Relationship:	Number:

HEALTHCARE PROVIDER INFORMATION

Practice Name:	Dr Name:	Phone:	Fax:
Address:		City:	State: Zip:
NPI:	DEA:	License:	UPIN:

INSURANCE INFORMATION *Provide Copy of Front & Back of Insurance Card *

Medical Insurance Name:	Prescription Insurance Name:
-------------------------	------------------------------

CLINICAL INFORMATION * Provide Medical Records *

TED: <input type="checkbox"/> Yes <input type="checkbox"/> No Diagnosis: <input type="checkbox"/> E05.00 <input type="checkbox"/> E06.3 <input type="checkbox"/> Other (specify): _____	Pregnancy Test: <input type="checkbox"/> Negative <input type="checkbox"/> Using Contraceptive Method
Other Comorbidities: _____ Weight: _____	
Therapy: <input type="checkbox"/> New to Therapy <input type="checkbox"/> Continuing Therapy <input type="checkbox"/> Next Due Date (if applicable): _____ <input type="checkbox"/> CAS Score (0-10) _____	
Laboratory Orders: <input type="checkbox"/> CBC <input type="checkbox"/> At Each Dose <input type="checkbox"/> Every _____ <input type="checkbox"/> CMP <input type="checkbox"/> At Each Dose <input type="checkbox"/> Every _____ <input type="checkbox"/> Other: _____	

PRESCRIPTION INFORMATION

Medication	Dose/Strength	Directions	Quantity	Refills
<input type="checkbox"/> Tepezza (teprotumumab-trbw)	<input type="checkbox"/> 500mg single dose vial	<input type="checkbox"/> Week 0: 10 mg/kg IV for 1 st infusion. Dose: _____ <input type="checkbox"/> Week 3: 20 mg/kg IV every 3 weeks for 7 infusions Dose: _____	<input type="checkbox"/> 21 day supply	

Reconstitute Tepezza vial with 10ml Sterile Water & gently swirl. **Dilute** with 0.9% Sod Chl. For doses < 1800mg use a 100ml bag & doses ≥ 1800mg use a 250mg bag. Discard 10.5ml of 0.9% Sodium Chloride from bag prior to adding diluted Tepezza into bag.

Administer the first 2 infusions over 90 minutes. Subsequent infusions may be reduced to 60 minutes if well tolerated. If infusion reaction occurs, interrupt or slow the rate of infusion.

Pre – Medication ** Not required but if patient experiences infusion reaction, consider premedicating 30 minutes prior infusion **

Medication	Dose	Directions	Refills	QTY
<input type="checkbox"/> Acetaminophen (Tylenol)	<input type="checkbox"/> 500mg <input type="checkbox"/> 1000mg PO			
<input type="checkbox"/> Diphenhydramine (Benadryl)	<input type="checkbox"/> 25mg <input type="checkbox"/> 50mg <input type="checkbox"/> PO <input type="checkbox"/> IV			
<input type="checkbox"/> Methylprednisolone (Solu-Medrol)	<input type="checkbox"/> 40mg <input type="checkbox"/> 125mg IV			
<input type="checkbox"/> 0.9% Sodium Chloride	<input type="checkbox"/> 5ml syringe <input type="checkbox"/> 10 ml syringe	Flush per protocol following infusion		
<input type="checkbox"/> Epinephrine Injection (Autoinjector)	<input type="checkbox"/> 0.3mg (Adult)			
<input type="checkbox"/> Anaphylaxis Medication Kit				
<input type="checkbox"/> Other:				

Nursing Coordination:
 YES: Provide home health nursing for education/training to administer reaction management 30 minute post-procedure observation.
 NO: Nursing not necessary: Patient and/or caregiver are proficient with infusion Physician to administer in the office
 Nursing already established, if so provide agency contact information: _____

PRESCRIBER SIGNATURE: _____	DATE: _____
Verbal Readback <input type="checkbox"/> Yes <input type="checkbox"/> No *Stamp signature not allowed, physician signature required *	

ADULT REACTION MANAGEMENT PROTOCOL

Observe for **hypersensitivity reaction**: Fever, chills, rigors, pruritus, rash, cough, sneezing, throat irritation, nausea, vomiting If reaction occurs:

- Stop infusion
- Maintain/establish vascular access
- Notify referring provider as clinically appropriate and follow clinical escalation protocol.
- EPSrx have the following PRN medications available for the following reactions.
 - Headache, pain, fever >100.4F, - Acetaminophen 500mg 1-2 tablets PO
 - Mild Hives, itching, redness, or rash - Diphenhydramine 25-50mg PO since dose, may repeat if needed
 - Severe hives, itching, redness, or rash- Diphenhydramine 25-50mg SIVP single dose, may repeat if needed
 - Nausea, vomiting, heartburn, acid reflux- Ondansetron 4mg ODT (may repeat x 1 in 20 minutes if nausea continues, max dose 8mg) or Famotidine 20mg PO
 - Hypotension (90/60), vasovagal response- Place patient in reclined position, administer 0.9% Sodium Chloride IV 250ml. May repeat to keep BP >90/60, maximum of 1000ml, monitor vital signs.
 - Chest pain/discomfort, shortness of breath- Call 911
 - Other _____
- When symptoms resolve resume infusion at 50% previous rate and increase per manufactures guidelines

Severe allergic/anaphylactic reaction:

- If symptoms are rapidly progressing or continuing after administration of prn medications above and signs symptoms of severe allergic/anaphylactic reaction (angioedema, swelling of the mouth, tongue, lips, or airway, dyspnea, bronchospasm with or without hypotension or hypertension.)
 - Call 911
 - Initiate basic life support as needed
 - **Epinephrine**- administer 0.3mg of a 1:1,000 (1mg/ml) concentration intramuscularly (preferably outer thigh), may be repeated every 5-15 minutes as needed to a maximum of 2 doses.
 - Place patient in recumbent position, elevate lower extremities
 - Administer **diphenhydramine** 50mg IV or Famotidine 20mg IVP, if not previously given
 - Administer **methylprednisolone** 125mg IVP, if not previously given.
 - Continuous monitoring of blood pressure, pulse oximetry, and heart rate.
 - Notify clinical executive, DON or CMO, when appropriate. Must be done same day. Do not delay treatment.

Patient Name

Patient Date of Birth

Provider Name (Print)

Provider Signature

Date