

**PATIENT INFORMATION**

Patient Name:		Date of Birth:	Gender:
Home Phone:	Cell Phone:	Email:	
Address:	City:	State:	Zip:
Emergency Contact:		Emergency Phone:	

**CLINICAL INFORMATION**

Patient Weight: \_\_\_\_\_  kg  lbs  
 Patient Height: \_\_\_\_\_  cm  in  
 Allergies: \_\_\_\_\_  
 Has patient previously received IG?  Yes  No  
 Patient already trained on subcutaneous infusion  
 Pharmacy to coordinate home health nursing visit and/or nursing training

**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_  
 DEA #: \_\_\_\_\_ NPI #: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_  
 Contact Person: \_\_\_\_\_

**PRESCRIPTION INFORMATION** Pharmacy to select IG if medication left blank

MEDICATION  Pharmacy to select IG  Brand \_\_\_\_\_

Infusion Route  Intravenous  Subcutaneous

**Dose and Directions**

➤ Pharmacy to calculate based on the following weight: \_\_\_\_\_  kg  lbs  
 Loading Dose: Infuse \_\_\_\_\_ GRAMS daily for \_\_\_\_\_ day(s)  
 -OR- Infuse \_\_\_\_\_ GRAM/KG divided over \_\_\_\_\_ day(s)  
 Maintenance Dose: Infuse \_\_\_\_\_ GRAMS daily for \_\_\_\_\_ days, every \_\_\_\_\_ week(s)  
 -OR- Infuse \_\_\_\_\_ GRAM/KG divided over \_\_\_\_\_ day(s), every \_\_\_\_\_ week(s)  
 Other \_\_\_\_\_

+/- 4 days to allow scheduling flexibility  OK to round to the nearest vial size  
 Multiple doses will be administered on consecutive days unless ordered otherwise.  
 Rate protocol: Titrate initial and maintenance infusions per manufacturer's product labeling.

**DIAGNOSIS**

G04.81 Other Encephalitis and Encephalomyelitis  
 M33.9 Dermatopolymyositis  
 D69.3 Immune Thrombocytopenia Purpura  
 M30.3 Kawasaki Disease  
 L12.9 Pemphigoid, Unspecified  
 L10.9 Pemphigus, Unspecified  
 M33.2 Polymyositis  
 Other \_\_\_\_\_

**REQUIRED FOR HOME INFUSION** RX to include diluents, needles, syringes, ancillary supplies, home medical equipment to administer infusion.

IV Access **To be administered PERIPHERALLY, unless otherwise indicated.**  PORT  PICC

**Flush Protocol** for IVIG drug admin days only  
 • 0.9% NaCl: 1-10mL IV before/after infusion, or PRN for line patency/SASH.  
 • Heparin 10 units/mL: 5mL IV (peripheral) PRN for final flush. \*For multi-day infusions  
 • Heparin 100 units/mL: 5mL IV (central) PRN for final flush.  
 \* For multi-day infusions, peripheral IV to be obtained, maintained, and discontinued for the duration of the infusion cycle per RN discretion in accordance with INS guidelines.

<b>Pre and Post Medications</b> Please strikethrough if not required	<b>To be given by mouth 30 minutes prior to infusion. May repeat every 4-6 hours as needed.</b>		<b>*For subcutaneous patients only* if requested by patient/nurse.</b>  <b>Lidocaine 2.5%/Prilocaine 2.5%</b> topical (may dispense Lidocaine 4%) to injection site(s) at least 1 hour prior to needle insertion.
	<b>Diphenhydramine</b>	25mg-50mg Adult max: 100mg/day	
	<b>Acetaminophen</b>	325mg-650mg Adult max: 3000mg/day	

**Anaphylaxis Protocol**  
**To be given intramuscularly PRN severe allergic reaction. Call 911. May repeat x 1.**  
 • Epinephrine 0.3mg (≥30kg/66lbs)

**Hydration**  
 Please select only if needed for IVIG  
 0.9% NaCl \_\_\_\_\_ mL infused over \_\_\_\_\_ minutes  
 D5W \_\_\_\_\_ mL infused over \_\_\_\_\_ minutes  
 Nurse to determine hydration rate if rate not provided above.  
**To be infused pre-infusion, unless otherwise indicated.**  
 Concurrent with Infusion  Other \_\_\_\_\_

**Diphenhydramine**  
 Please select only if needed for IVIG  
**To be given via slow IV push PRN for moderate – severe reaction.**  
 25-50mg  Other \_\_\_\_\_

**Quantity and Refills**  
 Dispense 1-month supply with 1-year refill unless indicated below.  
 Dispense 3-month supply with 1-year refill  Other \_\_\_\_\_

**Additional Orders**

**PRESCRIBER SIGNATURE REQUIRED (STAMP SIGNATURE NOT ALLOWED)**

By signing this form and using this pharmacy's services, you are authorizing this pharmacy to serve as your prior authorization designated agent in dealing with prescription and medical insurance companies.

<input type="checkbox"/> May Substitute/Product Selection Permitted/Substitution Permissible	<input type="checkbox"/> Dispense as Written/Brand Medically Necessary/Do Not Substitute/No Substitution/May Not Substitute
<b>Prescriber Signature</b> _____	<b>Prescriber Signature</b> _____
<b>Date</b> _____	<b>Date</b> _____

CA, MA, NC & PR: Interchange is mandated unless Prescriber writes the words "No Substitution" NY & Iowa providers, please submit electronic prescription.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to EPSrx or any of its subsidiaries using the contact information provided on this coversheet. This referral form is intended to assist with the processing of the patient's prescription and can be used at any pharmacy of the patient's choosing. It is not limited to the pharmacy listed on the form, nor does it imply that the patient is required to fill their prescription at a specific location. Patients maintain the right to select the pharmacy that best meets their needs. Please refer to TSBP Rule 281.7(15) for further guidance. EPS-CP/AI/7.1.2025